Author Index

Adair NE, 60
Adams J, 426
Agresti A, 307
Alison JA, 218
Angelopoulos TJ, 352
Arena R, 183

Bacher AC, 109, 115
Balza B, 139
Barefoot JC, 404
Barthélémy JC, 307
Bellin D, 100
Beohar N, 84
Berkebile ND, 260
Berkman LF, 398
Berry MJ, 60
Bittner V, 281
Black B, 29
Blackburn GG, 40
Blanchard CM, 29
Blumenthal J, 398
Braith RW, 201
Brawner CA, 260
Brolin SE, 228
Brown A, 17
Brubaker C, 352
Buchner D, 139
Burg MM, 404
Buzaitis A, 269
Bye PTP, 218

Cahill A, 17
Cannistra LB, 50
Carlson JJ, 322
Carney RM, 404
Castillo A, 426
Cecins NM, 228
Colansky MA, 40
Cooke D, 176
Coppersmith J, 139
Cornell CE, 404
Coronado M, 357
Costes F, 307
Courneya KS, 29
Court-Fortune I, 307
Czajkowski S, 398, 404

D'Angelo MS, 176
Dafoe W, 176, 442
Dasgupta H, 10
Daub B, 29
DeRegis JR, 109, 115
Després J, 161
DeVon HA, 122, 438
DiLillo V, 281

Ehrman JK, 260
Ettinger WH, 60

Falcone RA, 170
Ferrans CE, 122
Fesniak H, 10
Fitting J-W, 357
Fletcher D, 442
Fowler M, 100
Fraser SN, 29
Freedland KE, 404
Froelicher ES, 269, 398

Gardner JK, 299
Gerald L, 281
Gibbons LW, 266
Ginis KAM, 53
Ginter S, 269
Goble AJ, 415
Gordon NF, 349
Gosselink R, 143
Gullestad L, 100

Harris DE, 250
Hartman J, 426
Hasler CM, 334
Hauck CA, 299
Hawkes AL, 22
Haykowsky M, 208
Hees PS, 109
Herman CP, 299
Hevey D, 17
Hiatt WR, 170
Higginson L, 176
Hirsch AT, 170
Holt L, 139
Horgan JH, 17
Hosking JD, 404
Hsie M, 84
Huber M, 398
Humphrey R, 97, 183, 208
Hunziker J, 139

Ironson G, 398

Janssens J-P, 357
Jenkins SC, 228, 430
Johnson K, 176
Jordan S, 269

Kennedy MD, 208
Keteyian SJ, 260
Khatri P, 404
Kierns M, 17
Klinger TA, 299
Kundrat S, 334
Kwakkel G, 143

Lake FR, 430
Lakshminaryan S, 139
LaPier TK, 203
Larson EM, 314
Laubach Jr. CA, 299
Lauer MS, 88
Legro M, 139
Lichtenberger CM, 53
Lui K, 241, 365

MacKenzie CL, 53
Madigan MS, 183
Mahler DA, 226
Mandak JS, 10
McCartney N, 53
McConnell TR, 10, 190, 299
McKeough ZJ, 218
Merritt-Worden T, 151
Miller NH, 269
Misuraco A, 269
Mitchell PH, 398
Monti V, 322
Moore SM, 40
Morrin L, 153
Muralt B, 357
Murphy B, 415
Murray TC, 29
Myers J, 100

Newton H, 17
Norten J, 398
Nowak M, 22
Nuckolls J, 341

Oldridge N, 26, 438
Ouyang P, 109, 115

Pashkow FJ, 40
Peberdy MA, 183
Pelletier S, 334
Pfenninger P, 269
Phillips MM, 281
Pipe A, 176
Pitula CR, 398
Piña IL, 107
Poirier P, 161
Powell L, 398

Record NB, 250
Regensteiner JG, 170
Reid R, 176
Rejeski J, 60
Ries A, 78
Robinson E, 269
Roche F, 307
Rodgers WM, 29
Ross H, 100
Ruland CM, 40

Sanderson BK, 281
Schairer JR, 260
Schutz Y, 357
Segrest W, 426
Sevick MA, 60
Shapiro EP, 109
Sheps D, 404
Sherwood J, 269
Southard BH, 341
Southard DR, 341
Speare R, 22
Steele BG, 139
Stewart KJ, 109, 115, 170
Stojcevski Z, 415
Strauss GR, 430
Sung J, 109, 115
Sykes JS, 10

Takahashi T, 430
Tayback M, 109, 115
Taylor CB, 413
Taylor-Piliae RE, 90
Terrier PM, 357
Thomas RJ, 423
Tiukinhoy S, 84
Tokmakidis SP, 193
Tolentino R, 426
Treat-Jacobson D, 170
Turner KL, 109, 115

Vagelos R, 100
van't Hul A, 143
Vergnon J, 307
Vitcenda M, 290
Volaklis KA, 193

Wadley V, 269
Watson CP, 430
Williams MA, 170
Witta EL, 352
Worcester MUC, 415

Youngblood M, 398

Zaccaro DJ, 60

Subject Index

AACVPR. *See* American Association of Cardiovascular and Pulmonary Rehabilitation
Abdominal obesity, estimating, 109
Abstracts
  AACVPR 2003 Annual Meeting, 367
  CACR 13th Annual Meeting and Scientific Symposium, 387
Accelerometrycomparison with VO_2, during walk test, 384
  during respiratory rehabiltation, 357
  Activities of daily living, measures of, 378
  Adherence to exercise
  in COPD, 375
  factors associated with, 281
  theory of planned behavior and, 29
Aerobic exercise
  combined strength and aerobic exercise program, in CAD, 193
  frequency of, after cardiac rehabilitation, 371
  with Thera-Band® Elastic Bands, 388
African-Americans, 371
Agility measures, 378
American Association of Cardiovascular and Pulmonary Rehabilitation (AACVPR)
  2002-2003 AACVPR President, interview with, 151
  2003 Annual Meeting, 365
  2003 Annual Meeting abstracts, 367
  AACVPR News, 7, 151, 241, 385, 441
  award winners, 385
  International Heart and Lung Games, 386
Anthropometrics, 109
Arm exercise capacity, in COPD, 218
Automatic referral to cardiac rehabilitation, 387

Balance measures, 378
Baroreflex sensitivity, during exercise in heart failure secondary to dilated cardiomyopathy, 372
Behavior
  long-term outcomes after secondary prevention, 415
  planned, theory of, 29
Blood flow, aerobic training with Thera-Band® Elastic Bands and, 388
Blood glucose, elevated, lifestyle intervention in, 372
Blood lipids, combined-strength and aerobic exercise program and, in CAD, 193
Blood loss, in Talk Test, 384
Blood pressure, high, lifestyle intervention in, 373
Body image, and self-reported vs clinician-reported physiologic function, 53
Breathlessness, exercise training and, in heart failure, 10
Business model, quality patient care in, case report, 426

CACR. *See* Canadian Association of Cardiac Rehabilitation
CAD. *See* Coronary artery disease
Canadian Association of Cardiac Rehabilitation (CACR)
  13th Annual Meeting and Scientific Symposium abstracts, 387
  CACR News, 8, 153, 242, 442
  In-coming President's Report, 8
  Newsbeat, 153
Carbon dioxide
  VE/VCO_2 slope, prognostic value of, 371
  VE/VCO_2 slope vs VE/VCO_2 ratio, 370
Cardiac pacing, closed loop stimulation, 389
Cardiac rehabilitation
  ADLs after, measures of, 378
  aerobic exercise after, frequency of, 371
  aerobic training with Thera-Band® Elastic Bands for, 388
  in Caucasians vs African-Americans, 371
  in community settings, 250
  DANREHAB-study, 380
  with diabetes mellitus, 381
  exercise maintenance after, 369
  exercise maintenance after, predictors of, in women, 40
  failure to complete, factors associated with, 281
  four-week vs 10-week programs, 17
  functional status after, 379
  gender and diagnoses considerations for, 299
  HDL-cholesterol response to, 378
  and health care utilization and costs, 388
  and health outcomes, 388
  heart rate recovery after, 84
  maintenance, leisure time physical activity in, 260
  musculoskeletal pain and, 381
  nutrition education program for, 334
  in older adults, 382
  Ontario Cardiac Rehabilitation Pilot Project, final report, 242
  patient perceptions of magnitude of recovery during, 377
  phase II, exercise adherence during, 29
  quality of life after, 379
  quality of life during, 299
  risk stratification and, 388
  self-efficacy during, 299
  SF-36 Health Survey for, 22
  submaximal effort capacity during, 381
  tai chi in, 97
  twice daily, and surgical patient length of stay, 381
  with type 2 diabetes, 377
Cardiac rehabilitation programs
  case-managed home program, 387
  cost-effectiveness of standard vs alternative delivery of, 389
  geographic distribution in U.S., 378
  worship-based wellness programs, 374
Cardiac transplant recipients, physical activity patterns and exercise performance in, 100
Cardiology, preventive, measurement of improvements in outcomes, 423
Cardiomyopathy, dilated, heart failure secondary to, exercise in, 372
Cardiovascular risk
  resistance exercise training and, 201
  visceral obesity and, 161

waist circumference and, 161
Cardiovascular risk reduction programs
gender differences in, 376
INTERVENT program
educational status and, 376
in female vs male participants, 375
in premenopausal vs post-menopausal women, 376
in rural communities, 370
telephone/Internet-delivered, 370, 371
Cardiovascular screening, self-referred program for, 369
Case management
home programs, 387
Internet-based system, 341
Caucasians, 371
CHAMPS Physical Activity Questionnaire for Older Adults, 375
Cholesterol response to cardiac rehabilitation, 378
Chronic obstructive pulmonary disease
arm exercise capacity and dyspnea in, 218
constant-load cycle endurance performance in, 143
8-week outpatient pulmonary rehabilitation program in, 384
exercise adherence in, 375
heart rate and dyspnea in exercise prescription for, 228
long-term vs short-term exercise in, 60
noninvasive ventilation in exercise training in, 307
upper limb exercise test for, 430
Clinician-reported physiologic function, correlates of, 53
Closed loop stimulation cardiac pacing, effects on recovery heart rate and oxygen kinetics, 389
Combined-strength and aerobic exercise program, in CAD, 193
Community Healthy Activities Model Program for Seniors (CHAMPS) Physical Activity Questionnaire for Older Adults, 375
Community settings, cardiac rehabilitation in, 250
Congestive heart failure, hospital-based wellness program for, 352
Constant-load cycle endurance performance, in COPD, 143
COPD. *See* Chronic obstructive pulmonary disease
Coronary artery disease
combined-strength and aerobic exercise program in, 193
Enhancing Recovery in Coronary Heart Disease Trial (ENRICHD), 269
social support inventory, 398
functional status after hospitalization for, 203
risk with self-referred cardiovascular screening program, 369
smoking cessation in, 176
Cost-effectiveness of standard vs alternative service delivery, 389
Counseling, weekly, and weight loss, 368
Cycling
constant-load, in COPD, 143
music tempo and, 384

Daily activity, monitoring during pulmonary rehabilitation, 139
DANREHAB-study, 380
Daytime sleepiness, obstructive sleep apnea and, 377
Depression
aspects of social support associated with, 404
Enhancing Recovery in Coronary Heart Disease Trial (ENRICHD), 269
and self-reported vs clinician-reported physiologic function, 53
and smoking cessation, 368
Diabetes mellitus
cardiac rehabilitation with, 381
type 2
cardiac rehabilitation outcomes with, 377
exercise training and cardiovascular function in women with, 388
heart rate variability and isometric exercise response in obese women with, 372
Dialysis, massage and exercise for patients on, 383
Diaphragmatic function, after 8-week outpatient pulmonary rehabilitation program, in COPD, 384
Dietary patterns for secondary prevention, 322
Dilated cardiomyopathy, heart failure secondary to, exercise in, 372
Disability
and rehabilitative services for older adults after cardiac events, 380
resistance training effects in older women with, 379
Dual-energy X-ray absorptiometry, 109
Dyspnea
with COPD, 218
in exercise prescription with COPD, 228
after pulmonary rehabilitation, 367
target rate, 226
for exercise prescription in COPD, 228

Education, nutrition, 334
Educational status, and INTERVENT program effectiveness, 376
Endurance performance
constant-load cycle, in COPD, 143
measures of, after cardiac rehabilitation, 378
Enhancing Recovery in Coronary Heart Disease Trial (ENRICHD), 269
social support inventory, 398
Ethnic diversity, 441
Evaluation, SF-36 Health Survey, 22
Exercise
adherence to
in COPD, 375
factors associated with, 281
theory of planned behavior and, 29
aerobic, after cardiac rehabilitation, 371
arm capacity, in COPD, 218
capacity for
four-week vs 10-week programs and, 17
submaximal effort, during cardiac rehabilitation, 381
combined-strength and aerobic exercise program, in CAD, 193
endurance, constant-load cycle performance, in COPD, 143
intensity prescription with Talk Test, 382
isometric, hemodynamic response in type 2 obese diabetic women, 372
with left ventricular assist devices, 208
long-term vs short-term, in COPD, 60
maintenance
after cardiac rehabilitation, 369
women and, 50
with massage, for dialysis patients, 383
music tempo and, 384
performance in cardiac transplant recipients, 100
prescription for, in COPD
heart rate and dyspnea in, 228
target dyspnea rate (TDR) for, 228
target heart rate (THR) for, 228
tai chi, 90
tolerance for
exercise training and, in heart failure, 10
hospital-based wellness program for maintenance of, in CHF, 352
noninvasive ventilation and, in COPD, 307
and weight loss, 368
women's maintenance after cardiac rehabilitation, predictors of, 40
Exercise-induced glutathione oxidative stress, max GXT assessment of, 383

Exercise tests
6-minute walk vs NuStep, 383
upper limb, in COPD, 430
Exercise training
and cardiovascular disease risk, 201
and cardiovascular function in women with type 2 diabetes mellitus, 388
for heart failure, 10
with left ventricular assist devices, 208
monitoring, Session RPE method for, 382
noninvasive ventilation during, in COPD, 307
and oxidative stress, 373
External counterpulsation treatment, quality of life after, 376

Faithfully Fit Forever (426), 374
Fasting blood glucose, elevated, lifestyle intervention in, 372
Fatness, and health-related quality of life in older persons, 115
Fitness, and health-related quality of life in older persons, 115
Flexibility, measures of, after cardiac rehabilitation, 378
Foods, functional, 334
Functional status
after cardiac rehabilitation, 379
after hospitalization for CHD, 203
physical activity and, in adult women, 383

Gender differences
in comprehensive cardiovascular risk reduction programs, 375
in HDL-cholesterol response to cardiac rehabilitation, 378
in intensive comprehensive risk factor reduction program, 376
in quality of life and self-efficacy, 299
Geographic distribution of CR programs in U.S., 378
Glucose
abnormal metabolism of, 380
elevated fasting blood, lifestyle intervention in, 372
Glutathione oxidative stress, exercise-induced, max GXT assessment of, 383

Health care utilization and costs, 388
Health-related quality of life, in older men and women, 115
Heart disease
activity limitations and, 378
coronary
Enhancing Recovery in Coronary Heart Disease Trial (ENRICHD), 269, 398
functional status after hospitalization for, 203
risk with self-referred cardiovascular screening program, 369
exercise training and, in women with type 2 diabetes, 388
Internet-based case management system for, 341
MacNew Heart Disease Health-related Quality of Life questionnaire, 122, 438
secondary prevention of
dietary patterns for, 322
Internet-based case management system for, 341
Heart failure
congestive, hospital-based wellness program for, 352
exercise training for, 10
peak oxygen consumption in, predicting, 183
prognostic value of VE/VCO_2 slope in, 371
prognostic value of VE/VCO_2 slope vs VE/VCO_2 ratio in, 370
secondary to dilated cardiomyopathy, 372
"Heart healthy" functional foods, 334
Heart rate
in exercise prescription with COPD, 228
recovery
after cardiac rehabilitation, 84
closed loop stimulation cardiac pacing and, 389
as modifiable risk factors, 88
response to exercise in heart failure secondary to dilated cardiomyopathy, 372
target, 226
for exercise prescription in COPD, 228
variability in type 2 obese diabetic women, 372
High-density lipoprotein–cholesterol response to cardiac rehabilitation, 378
Home programs, case-managed, 387
Hospital-based wellness program, for CHF, 352
Hospitalization
depression at, aspects of social support associated with, 404
functional status after, 203
smoking cessation in, 176
surgical patient length of stay, twice daily cardiac rehabilitation and, 381
Hypertension, lifestyle intervention in, 373

International Heart and Lung Games, 386
Internet
-based case management system for secondary prevention, 341
-delivered INTERVENT program
clinical effectiveness of, 371
in rural communities, 370
service delivery via, 349
WSCVPR Web-based Outcomes Project (W^2eBOP), 290
INTERVENT Lifestyle Management and Cardiovascular Risk Reduction Program (INTERVENT)
educational status and, 376
in female vs male participants, 375
in premenopausal vs post-menopausal women, 376
telephone/Internet-delivered
clinical effectiveness of, 371
in rural communities, 370
Interview administration of CHAMPS Physical Activity Questionnaire for Older Adults, 375
Isolation, social, 269

Left ventricular assist devices, 208
Leisure time physical activity, 260
Length of stay, surgical patient, twice daily cardiac rehabilitation and, 381
Lifestyle intervention
in elevated fasting blood glucose, 372
gender differences in, 376
in hypertension, 373
intensive comprehensive risk factor reduction program for, 376
INTERVENT program
educational status and, 376
in female vs male participants, 375
in premenopausal vs post-menopausal women, 376
in rural communities, 370
telephone/Internet-delivered, 370, 371
Lipids, blood, combined-strength and aerobic exercise program and, in CAD, 193
Literature Update, 69, 235
Lung transplant recipients, resistance training for, 367

MacNew Heart Disease Health-related Quality of Life questionnaire, 122, 438
Magnetic resonance imaging of abdominal obesity, 109
Maintenance cardiac rehabilitation, leisure time physical activity in, 260
Massage, for dialysis patients, 383
Men
comprehensive cardiovascular risk reduction program for, 375
older

health-related quality of life and mood in, 115
MRI of abdominal obesity in, 109
Minute ventilation (VE)
VE/VCO_2 slope, prognostic value of, 371
VE/VCO_2 slope vs VE/VCO_2 ratio, 370
Modified Bruce Protocol, 383
Monitoring daily activity, 139
Monitoring exercise training, Session RPE method for, 382
Mood
obstructive sleep apnea and, 377
in older men and women, 115
MRI. *See* Magnetic resonance imaging
Multidisciplinary cardiac rehabilitation, four-week vs 10-week programs, 17
Multidisciplinary risk reduction model, first results, 369
Muscular strength, Thera-Band® Elastic Bands and, 388
Musculoskeletal pain, and cardiac rehabilitation, 381
Music tempo, and cycling performance, 384
Myocardial infarction
cardiac rehabilitation for, 299
Enhancing Recovery in Coronary Heart Disease Trial (ENRICHD), 269, 398
Quality of Life After Myocardial Infarction Questionnaire (QLMI), 122, 438
social support for patients after, 413

National Heart, Lung, and Blood Institute (NHLBI) workshop on peripheral arterial disease, 241
Newsbeat (CACR), 153
Nicotine dependence, 368
NIDDM. *See* Diabetes, type 2
North Carolina pulmonary rehabilitation programs
dyspnea after participation in, 367
quality of life after participation in, 367
NuStep test, 383
Nutrition
dietary patterns for secondary prevention, 322
education program for cardiac rehabilitation, 334

Obesity
abdominal, estimating, 109
heart rate variability and isometric exercise response in type 2 diabetic women with, 372
visceral, and cardiovascular risk, 161
Obstructive sleep apnea, and daytime sleepiness, mood, and quality of life, 377
Old old, typology of, based on SF-36, 379
Older adults
CHAMPS Physical Activity Questionnaire for Older Adults, 375
disability and rehabilitative services after cardiac events, 380
disabled women, resistance training effects on, 379
health-related quality of life and mood in, 115
MRI of abdominal obesity in, 109
physical function and, 382
response to cardiac rehabilitation, 382
Ontario Cardiac Rehabilitation Pilot Project, final report, 242
Osteoporosis, vertebral, resistance training for prevention of, 367
Outpatient pulmonary rehabilitation programs
8-week, in COPD, 384
at tertiary care centers, 368
Oxidative stress
exercise-induced glutathione, max GXT assessment of, 383
exercise training and, 373
Oxygen consumption (VO_2)
comparison with accelerometry, during walk test, 384
peak, predicting, 183
Oxygen kinetics, closed loop stimulation cardiac pacing and, 389

Pacing, closed loop stimulation, 389
Pain, musculoskeletal, 381
Patient compliance, factors associated with, 281
Patient education, nutrition program, 334
Patient perceptions
of magnitude of recovery during cardiac rehabilitation, 377
of quality of life, 379
Percutaneous coronary intervention, 299
Peripheral arterial disease
NHLBI workshop or:, 241
rehabilitation for, 170
Phase II cardiac rehabilitation, theory of planned behavior and exercise adherence during, 29
Physical activity
CHAMPS Physical Activity Questionnaire for Older Adults, 375
commitment to, 107
daily, monitoring during pulmonary rehabilitation, 139
effects on functional performance in adult women, 383
and health-related quality of life in older persons, 115
leisure time, in maintenance cardiac rehabilitation, 260
limitations on, and heart disease, 378
patterns in cardiac transplant recipients, 100
walking activity, by accelerometry, 357
Physical activity limitations
and heart disease, 378
Physical function
and older adults, 382
resistance training and, in disabled older females, 379
self-reported vs clinician-reported, correlates of, 53
Planned behavior, theory of, 29
Prevention, secondary
dietary patterns for, 322
Internet-based case management system for, 341
long-term behavioral outcomes, 415
service delivery in, 349
Preventive cardiology, measurement of improvements in outcomes, 423
Psychosocial Risk Factor Survey, evaluation of, 374
Pulmonary disease. *See* Chronic obstructive pulmonary disease
Pulmonary rehabilitation
dyspnea after, 367
monitoring daily activity during, 139
outpatient programs
8-week, in COPD, 384
at tertiary care centers, 368
perspectives on, 78
quality of life after, 367

Q-factor analysis, for typology of old old, 379
Quality of life
after cardiac rehabilitation, 379
after exercise training, in heart failure, 10
after external counterpulsation treatment, 376
four-week vs 10-week programs and, 17
gender differences and diagnoses considerations for, 299
health-related, in older men and women, 115
hospital-based wellness program for maintenance of, in CHF, 352
instruments, psychometric properties of, 122, 438
obstructive sleep apnea and, 377
patient vs proxy perceptions of, 379
after pulmonary rehabilitation, 367
Quality of Life After Myocardial Infarction Questionnaire (QLMI), 122, 438
Quality of Life Index–Cardiac Version 441 (QLI-441), 122
Quality patient care, in business model, case report, 426

Racial/ethnic diversity, 441
Recovery during cardiac rehabilitation, patient perceptions of, 377
Referral to cardiac rehabilitation, automatic, 387
Rehabilitation. *See also* Cardiac rehabilitation; Pulmonary rehabilitation
  factors associated with adherence to, 281
  for older adults after cardiac events, 380
  peripheral arterial disease, 170
  respiratory, accelerometry during, 357
Rehabilitation programs
  four-week vs 10-week, 17
  racial/ethnic diversity in, 441
  SF-36 Health Survey for, 22
Resistance training
  and cardiovascular disease risk, 201
  and physical performance in disabled older females, 379
  for prevention of vertebral osteoporosis, 367
Respiratory muscle endurance, exercise training and, in heart failure, 10
Respiratory rehabilitation, accelerometry during, 357
Risk reduction
  multidisciplinary model for, first results, 369
  telephone/Internet-delivered program in rural communities, 370
Risk stratification, 388
Rockport Fitness Walking Test (RFWT), 384
Rural communities, telephone/Internet-delivered INTERVENT program in, 370, 371

Screening, self-referred program for, 369
Seattle Angina Questionnaire (SAQ), 122
Secondary prevention
  dietary patterns for, 322
  Internet-based case management system for, 341
  long-term behavioral outcomes, 415
  service delivery in, 349
Self-administration of CHAMPS Physical Activity Questionnaire for Older Adults, 375
Self-efficacy, gender differences and diagnoses considerations for, 299
Self-referred cardiovascular screening program, 369
Self-regulation, and exercise adherence in COPD, 375
Self-reported physiologic function, correlates of, 53
Senior Fitness Test, 378
Service delivery
  cost-effectiveness of standard vs alternative, 389
  via Internet, 349
Session RPE method, for monitoring exercise training, 382
Short Form-36 Health Survey (SF-36), 22
  patient vs proxy responses to, 379
  psychometric properties of, 122
  typology of old old based on, 379
Sibutramine, for weight loss after smoking cessation, 374
6-minute walk test, 383
Sleep apnea, obstructive, and daytime sleepiness, mood, and quality of life, 377
Sleepiness, daytime, obstructive sleep apnea and, 377
Smoking cessation
  in CAD, 176
  depression and, 368
  sibutramine for weight gain after, 374
  stepped care approach to, 176
Social support, 413
  aspects of, 404
  Enhancing Recovery in Coronary Heart Disease Trial (ENRICHD), 269
  ENRICHD inventory, 398
Stepped care approach to smoking cessation, 176
Strength measures, after cardiac rehabilitation, 378
Strength training, combined with aerobic exercise, in CAD, 193
Stress, oxidative
  exercise-induced glutathione, max GXT assessment of, 383
  exercise training and, 373
Surgical patient length of stay, twice daily cardiac rehabilitation and, 381
Surgical revascularization, cardiac rehabilitation after, 299

Tai chi, 90, 97
Talk Test
  blood loss and ventilatory threshold in, 384
  exercise intensity prescription with, 382
Target dyspnea rate (TDR), 226
  in COPD, 228
Target heart rate (THR), 226
  in COPD, 228
TDR. *See* Target dyspnea rate
Telephone/Internet-delivered INTERVENT program
  clinical effectiveness of, 371
  in rural communities, 370
Tertiary care centers, outpatient pulmonary rehabilitation program in, 368
Thera-Band® Elastic Bands, 388
THR. *See* Target heart rate
Tobacco use. *See* Smoking
Training
  aerobic, with Thera-Band® Elastic Bands, 388
  combined-strength and aerobic exercise program, in CAD, 193
  exercise
    and cardiovascular disease risk, 201
    and cardiovascular function in women with type 2 diabetes mellitus, 388
    for heart failure, 10
    with left ventricular assist devices, 208
    monitoring, Session RPE method for, 382
    noninvasive ventilation during, in COPD, 307
    and oxidative stress, 373
  resistance
    and cardiovascular disease risk, 201
    and physical performance in disabled older females, 379
    for prevention of vertebral osteoporosis, 367
Transplantation
  physical activity patterns and exercise performance in cardiac transplant recipients, 100
  resistance training for lung transplant recipients, 367
Treadmill GXT, maximal, for assessment of exercise-induced glutathione oxidative stress, 383

United States, geographic distribution of CR programs in, 378
Upper limb exercise test, for COPD, 430

Ventilation
  minute (VE)
    VE/VCO_2 slope, prognostic value of, 371
    VE/VCO_2 slope vs VE/VCO_2 ratio, 370
  noninvasive, in COPD, 307
Ventilatory threshold, in Talk Test, 384
Vertebral osteoporosis, resistance training for prevention of, 367
Visceral obesity, and cardiovascular risk, 161
Waist circumference, and cardiovascular risk, 161
Walking test
  accelerometry, during respiratory rehabilitation, 357
  accelerometry vs VO_2 during, 384
Weekly counseling, and weight loss, 368

Weight loss
sibutramine for, after smoking cessation, 374
weekly counseling and exercise effects on, 368
Wellness programs
hospital-based, for CHF, 352
worship-based, 374
Wisconsin Society for Cardiovascular and Pulmonary Rehabilitation (WSCVPR) Web-based Outcomes Project (W^2eBOP), 290
Women
exercise maintenance for, 50
INTERVENT program for, 375, 376
obese, with type 2 diabetes, heart rate variability and isometric exercise response in, 372
older
with disabilities, resistance training effects on, 379
health-related quality of life and mood in, 115
MRI of abdominal obesity in, 109
physical activity and functional performance in, 383
predictors of exercise maintenance for, 40
with type 2 diabetes
exercise training and cardiovascular function in, 388
obese, heart rate variability and isometric exercise response in, 372
Worship-based wellness programs, 374
WSCVPR Web-based Outcomes Project (W^2eBOP), 290

X-ray absorptiometry, dual-energy, 109

STATEMENT OF OWNERSHIP, MANAGEMENT, AND CIRCULATION (Act of August 12, 1970; Section 3685; Title 39 United States Code) Date of Filing—October 1, 2003. Title of Publication—*Journal of Cardiopulmonary Rehabilitation*; Frequency of Issue—Bimonthly; Annual Subscription Price—$97.00; Location of Known Office of Publication—Lippincott Williams & Wilkins, Inc., 16522 Hunters Green Parkway, Hagerstown, MD 21740-2116; Location of the Headquarters or General Business Offices of the Publisher—Lippincott Williams & Wilkins, Inc., 530 Walnut Street, Philadelphia, PA 19106; Publisher—Lippincott Williams & Wilkins, Inc., 530 Walnut Street, Philadelphia, PA 19106; Editor—Gary Balady, MD, Boston University Medical Center, Preventive Cardiology, 88 East Newton Street, Boston, MA 02118; Managing Editor—Dana Fennewald, American Association of Cardiovascular and Pulmonary Rehabilitation, 401 North Michigan Avenue, Suite 2200, Chicago, IL 60611-4267; Owner—Lippincott Williams & Wilkins, 530 Walnut Street, Philadelphia, PA 19106; 351 West Camden Street, Baltimore, MD 21201; Wolters Kluwer, US, 345 Hudson Street, New York, NY 10014; Wolters Kluwer nv (owns 100% of stock), Stadouderskade 1, 1054 FS Amsterdam, The Netherlands; Known Bond Holders, Mortgagees, and other security holders owning or holding 1 percent or more of the total amount of bonds, mortgages, or other securities—None. A. Total no. of copies printed (net press run), average 4,875, actual 5,200. B. Paid and/or requested circulation 1. Paid/requested outside-county mail subscriptions stated on form 3541, average 3,990, actual 3,661; 2. Paid in-county subscriptions, none; 3. Sales through dealers and carriers, street vendors, counter sales, and other non-USPS paid distribution, average 661, actual 676; 4. Other classes mailed through the USPS, none. C. Total paid and/or requested circulation [sum of B (1), (2), (3), and (4)], average 4,651, actual 4,337. D. Free distribution by mail (samples, complimentary, and other free), 1. Outside-county as stated on form 3541, average 146, actual 188; 2. In-county as stated on form 3541, none; 3. Other classes mailed through the USPS, none. E. Free distribution outside the mail (carriers or other means), average 34, actual 52. F. Total free distribution (sum of D and E), average 180, actual 240. G. Total distribution (sum of C and F), average 4,831, actual 4,577. H. Copies not distributed, average 705, actual 1,299. I. Total (sum of G and H), average 5,536, actual 5,876. Percent paid and/or requested circulation, average 96.27%, actual 94.76%. I certify that the statements made by me above are correct and complete. *Jeffrey Brown, Manager, Periodicals Operations.*